


PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen



Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

Ansökan ingavs ursprungligen på engelska.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

The application was originally filed in English.

(71) Sökande AstraZeneca AB, Södertälje SE
Applicant (s)

(21) Patentansökningsnummer 0302116-9
Patent application number

(86) Ingivningsdatum 2003-07-21
Date of filing

Stockholm, 2004-07-09

För Patent- och registreringsverket
For the Patent- and Registration Office

Hjordis Segerlund

Avgift
Fee 170:-

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

PATENT- OCH
REGISTRERINGSVERKET
SWEDEN

Postadress/Adress
Box 5055
S-102 42 STOCKHOLM

Telefon/Phone
+46 8 782 25 00
Vx 08-782 25 00

Telex
17978
PATOREG S

Telefax
+46 8 666 02 86
08-666 02 86

NOT FOR PUBLICATION

NOVEL COMPOUNDS

- 5 The present invention relates to piperidine compounds, processes and intermediates used in their preparation, pharmaceutical compositions containing them and their use in therapy.

Histamine is a biogenic amine that regulates a variety of physiological and pathological processes including inflammation, gastric acid secretion and neurotransmission.


- 10 Histamine acts via a family of G-protein coupled receptors and 4 members of this family have been identified and cloned: histamine H1 (Yamashita *et al*, 1991), histamine H2 (Gantz *et al*, 1991), histamine H3 (Lovenberg *et al*, 1991) and histamine H4 receptor (Oda *et al*, 1999). H1 and H2 are the best characterised of these receptors and antagonists of both are used clinically. In general, inflammatory and allergic responses are modified by
- 15 H1 receptors (Ash and Schild, 1966) while gastric acid secretion is regulated by interaction with H2 receptors (Black *et al*, 1972). H1 antagonists are therefore used to treat a variety of allergic conditions and H2 antagonists are used to treat gastric ulcers. Histamine appears to regulate neurotransmitter release via H3 receptors (Arrang *et al*, 1983), but the role of the recently identified histamine H4 receptor is currently unknown. The histamine
- 20 H4 receptor bears sequence and pharmacological similarity to the H3 receptor, although the tissue distribution profiles of both receptors are different. The H3 receptor is abundant in the brain and neural tissue while the H4 receptor appears to be restricted to peripheral tissues. The H4 receptor has a high distribution in peripheral blood leukocytes, especially eosinophils and neutrophils and H4 mRNA expression has also been demonstrated in other
- 25 immune and inflammatory cells, including T-cells, dendritic cells, monocytes, macrophages, mast cells and epithelial cells. In addition, there is some evidence that receptor expression may be modulated by cytokine activation (Morse *et al*, 2001). The H4 receptor may therefore have a role in immune and/or inflammatory modulation.

- 30 Histamine H1 receptor antagonists are successfully used in the treatment of allergic rhinitis but provide incomplete blockade of all symptoms resulting in the need for co-administration of other agents to treat nasal congestion, usually sympathomimetic amine decongestants. Combinations of H1 and H2 antagonists also fail to give complete blockade of these effects. Similarly, although histamine contributes to many of the

physiological processes that occur in asthma, histamine H1 antagonists are not used in asthma because of inconsistent efficacy. Some further anti-inflammatory activity appears to be required to block the effects of histamine in many patho-physiological processes, implying a role for additional pro-inflammatory histamine receptors. The H4 receptor may serve such a role, and agents that interact with H4 receptors either alone, or in combination with other histamine receptors or anti-inflammatory agents, may provide enhanced efficacy in disease.

Antagonists of the histamine H4 receptor may therefore have utility in a variety of diseases or disorders. WO 02/072548 discloses a series of compounds said to be active as mediators of the histamine H4 receptor.

There is a need for further compounds having this activity, and the present invention therefore provides a compound of formula (I) and pharmaceutically acceptable salts and solvates thereof for use in the manufacture of a medicament for the treatment of diseases mediated by histamine H3 and H4:



in which:

Ar is an aryl group, a 5-7 membered heteraromatic ring containing 1-4 heteroatoms selected from nitrogen, oxygen or sulphur, or a bicyclic or tricyclic heteraromatic ring containing 1-4 heteroatoms selected from nitrogen, oxygen or sulphur, each of which can be optionally substituted by 1-3 groups selected from C₁₋₆ alkyl, C₁₋₆ alkylthio, C₁₋₆ alkoxy, halogen, cyano, CF₃, OCF₃, C₃₋₆ cyclolalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₂₋₆ alkenyloxy, hydroxyl, nitro, tosyl, thienyl, benzyl, phenyl, nitrophenyl,

R¹ is hydrogen or C₁₋₆ alkyl;

X is O, NR^2 , CH_2 or SO_x

R^2 is C_{1-6} alkyl;

5

x is 0, 1 or 2;

Y is CH_2 , $\text{C}=\text{O}$, SO_2 , or $(\text{C}=\text{O})\text{NH}$;

10 Z is $(\text{CR}^3\text{R}^4)_r$ or Y and Z together form a $\text{CH}=\text{CH}$ group;

m and n are independently 0, 1, 2 or 3;

p and q are independently 0, 1 or 2;

15

r is 0, 1, 2, 3, or 4 and

R^3 and R^4 are independently hydrogen or C_{1-6} alkyl.

20 Certain compounds of formula (I) are capable of existing in stereoisomeric forms. It will be understood that the invention encompasses all geometric and optical isomers of the compounds of formula (I) and mixtures thereof including racemates. Tautomers and mixtures thereof also form an aspect of the present invention.

25 The term aryl includes phenyl and naphthyl. The term alkyl, whether alone or as part of another group, includes straight chain and branched chain alkyl groups. Examples of 5- to 7-membered heteroaromatic ring containing 1 to 4 heteroatoms include thienyl, furanyl, pyrrolyl, imidazolyl, pyridyl, pyrazinyl, pyrimidyl, pyridazinyl, triazinyl, oxazolyl, thiazolyl, isoxazolyl, pyrazolyl, oxadiazolyl, thiadiazolyl, triazolyl and tetrazolyl..

30 Examples of suitable bicyclic rings include indole, benzothiophene, quinoline, benzodioxan, and naphthyl. Examples of suitable tricyclic rings include dibenzofuran and thiene[2,3-b]benzothiophene. For any of these mono- bicyclic or tricyclic rings, substituents can be present in any suitable ring position including suitable substituents on nitrogen atoms.

35 Preferably Ar^1 is phenyl, furyl or thienyl optionally substituted as defined above. More preferably Ar^1 is phenyl optionally substituted as defined above. Preferred substituents

include halogen such as iodo, chloro and fluoro, cyclohexyl, methyl, ethyl, propyl, t-butyl, ethynyl, propenyloxy, hydroxyl, methoxy, nitro, tosyl, trifluoromethyl, thienyl, benzyl, cyano, phenylethynyl, nitrophenyl, methylthio, propoxy, butoxy, 2-propenyl, or trifluomethoxy.

5

Most preferably Ar¹ is phenyl substituted by bromo, hydroxyl or 2,4-difluoro.

Preferably R¹ is hydrogen or methyl.

10

Preferably X is O.

Preferably Y is CH₂ or C=O and Z is CH₂, CHMe, CH₂CHMe or Y and Z form a CH=CH group.

15

More preferably Y is CH₂ or C=O and Z is CH₂.

Preferably m is 1 and n is 0.

Preferably p and q are both 1.

20

Preferred compounds of the invention include:

4-(1*H*-Imidazol-4-ylmethoxy)-1-(1-oxo-3-phenylbutyl)-piperidine

4-(1*H*-Imidazol-4-ylmethoxy)-1-[[4-(trifluoromethyl)phenyl]acetyl]-piperidine

1-[2-(4-Hydroxyphenyl)-1-oxopropyl]-4-[(5-methyl-1*H*-imidazol-4-yl)methoxy]-

25

piperidine

1-[(4-fluorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine

1-[(2-chlorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine

1-[(4-chlorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine

4-(1*H*-imidazol-4-ylmethoxy)-1-(phenylacetyl)-piperidine

30

1-(4-cyclohexylbenzoyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine

1-[(3,4-dichlorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine

4-(1*H*-imidazol-4-ylmethoxy)-1-[(4-methylphenyl)acetyl]-piperidine

1-[(3,4-difluorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine


- 1-[(2,4-difluorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(4'-propyl[1,1'-biphenyl]-4-yl)carbonyl]-piperidine
1-[2-(4-hydroxyphenyl)-1-oxopropyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(2*E*)-3-(3,4-dichlorophenyl)-1-oxo-2-propenyl]-4-(1*H*-imidazol-4-ylmethoxy)-
5 piperidine
1-[3-(2,4-dichlorophenyl)-1-oxopropyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(2,4-dichlorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(2-Bromophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(3-Bromo-2-thienyl)methyl]-4-[(5-methyl-1*H*-imidazol-4-yl)methoxy]-piperidine
10 1-[(3-bromo-2-thienyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(4-ethynylphenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[[3-(4-methylphenoxy)phenyl]methyl]-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[[4-(2-propenyloxy)phenyl]methyl]-piperidine
4-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-phenol
15 4-(1*H*-imidazol-4-ylmethoxy)-1-[(2-methoxyphenyl)methyl]-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[[3-(4-methoxyphenoxy)phenyl]methyl]-piperidine
1-[(2,3-dichlorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(2-chloro-4-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-(2-dibenzofuranylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
20 4-(1*H*-imidazol-4-ylmethoxy)-1-[[2-(methylthio)phenyl]methyl]-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-(thieno[2,3-*b*][1]benzothien-2-ylmethyl)-piperidine
1-[(2-chloro-5-nitrophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1*H*-pyrrole, 2-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-1-[(4-
methylphenyl)sulfonyl]-
25 2-ethoxy-6-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-phenol
1-(1,3-benzodioxol-5-ylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[[4-(phenylmethoxy)phenyl]methyl]-piperidine
1-[[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(4-bromophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
30 4-(1*H*-imidazol-4-ylmethoxy)-1-[(4-methylphenyl)methyl]-piperidine

- 4-(1*H*-imidazol-4-ylmethoxy)-1-(2-thienylmethyl)-piperidine
1-[(4-chlorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(2-chloro-6-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(3-methyl-2-thienyl)methyl]-piperidine
5 4-(1*H*-imidazol-4-ylmethoxy)-1-(2-naphthalenylmethyl)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-(1-naphthalenylmethyl)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(2-nitrophenyl)methyl]-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-(3-thienylmethyl)-piperidine
1-[(1,1'-biphenyl)-4-ylmethyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
10 1-[(2,5-difluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(3-phenoxyphenyl)methyl]-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(3-methylphenyl)methyl]-piperidine
1-(2-furanyl)methyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(2,6-dichlorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
15 1-[(4-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(3-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-(3-furanyl)methyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(4-ethylphenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(2-methylphenyl)methyl]-piperidine
20 1-[(3-chlorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(5-methyl-2-thienyl)methyl]-piperidine
1-[(4-bromo-2-thienyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(2,2'-bithiophen)-5-ylmethyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
3,5-dichloro-2-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-phenol
25 1-[(3,4-difluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(3,5-difluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(6-chloro-1,3-benzodioxol-5-yl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[[4-[4-(1,1-dimethylethyl)-2-thiazolyl]phenyl]methyl]-4-(1*H*-imidazol-4-ylmethoxy)-
piperidine
30 4-(1*H*-imidazol-4-ylmethoxy)-1-[(1-methyl-1*H*-pyrrol-2-yl)methyl]-piperidine

- 1*H*-indole, 3-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-1-(phenylmethyl)-
1-[(5-chloro-2-thienyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-(1,3-benzodioxol-4-ylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
2-thiophenecarbonitrile, 3-[[4-[[4-(1*H*-imidazol-4-ylmethoxy)-1-
5 piperidinyl]methyl]phenoxy]methyl]-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[[5-(phenylethynyl)-2-thienyl]methyl]-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[[5-(4-nitrophenyl)-2-furanyl]methyl]-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[[5-(3-nitrophenyl)-2-furanyl]methyl]-piperidine
1-[(4-chloro-1*H*-pyrazol-3-yl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
10 1-[(4-bromo-1-methyl-1*H*-pyrazol-3-yl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(4-bromo-1*H*-pyrazol-3-yl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
2-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-benzonitrile
4-(1*H*-imidazol-4-ylmethoxy)-1-[(4-iodophenyl)methyl]-piperidine
1-[(5-ethyl-2-thienyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
15 4-(1*H*-imidazol-4-ylmethoxy)-1-[[5-(methylthio)-2-thienyl]methyl]-piperidine
1-[[1-(3,5-dichlorophenyl)-1*H*-pyrrol-2-yl]methyl]-4-(1*H*-imidazol-4-ylmethoxy)-
piperidine
1-[[1-(4-chlorophenyl)-1*H*-pyrrol-2-yl]methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[[4-(phenylethynyl)-2-thienyl]methyl]-piperidine
20 4-(1*H*-imidazol-4-ylmethoxy)-1-[(3-phenoxy-2-thienyl)methyl]-piperidine
1-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(4-propoxyphenyl)methyl]-piperidine
2-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-phenol
1-[(2,4-difluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
25 3-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-2-thiophenecarbonitrile
1-(benzo[*b*]thien-3-ylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
2-chloro-3-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-pyridine
3-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-2-(2-propenyl)-phenol
1-[(4-chloro-3-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
30 4-(1*H*-imidazol-4-ylmethoxy)-1-[[4-(trifluoromethoxy)phenyl]methyl]-piperidine

1-[(2,6-difluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-[(4-bromo-2-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-[(2,2-difluoro-1,3-benzodioxol-5-yl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-[(4-butoxyphenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 4-(1*H*-imidazol-4-ylmethoxy)-1-[(2,3,5-trichlorophenyl)methyl]-piperidine
 1-[(2,5-dichlorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 4-(1*H*-imidazol-4-ylmethoxy)-1-[[2-(trifluoromethyl)phenyl]methyl]-piperidine
 1-[(4-chloro-2-nitrophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 and pharmaceutically acceptable salts and solvates thereof.

Certain compounds of formula (I) are novel, and all novel compounds of formula (I) form a further aspect of the invention. The invention therefore provides a compound of formula (IA):



(IA)

in which:

Ar is an aryl group, a 5-7 membered heteraromatic ring containing 1-4 heteroatoms selected from nitrogen, oxygen or sulphur, or a bicyclic or tricyclic heteraromatic ring containing 1-4 heteroatoms selected from nitrogen, oxygen or sulphur, each of which can be optionally substituted by 1-3 groups selected from C₁₋₆ alkyl, C₁₋₆ alkylthio, C₁₋₆ alkoxy, halogen, cyano, CF₃, OCF₃, C₃₋₆ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₂₋₆ alkenyloxy, hydroxyl, nitro, tosyl, thienyl, benzyl, phenyl, nitrophenyl,

R¹ is hydrogen or C₁₋₆ alkyl;

X is O, NR², CH₂ or SO_x

R² is C₁₋₆ alkyl;

x is 0, 1 or 2;

Y is C=O, SO₂, or (C=O)NH;

5

Z is (CR³R⁴)_r or Y and Z together form a CH=CH group;

m and n are independently 0, 1, 2 or 3;

10

p and q are independently 0, 1 or 2;

r is 0, 1, 2, 3, or 4, and

R³ and R⁴ are independently hydrogen or C₁₋₆alkyl.


15

For compounds (IA) Y is preferably C=O. Other preferred substituents for compounds of formula (IA) are those defined above.

20

According to the invention there is also provided a process for the preparation of compounds (I)/(IA) which comprises:


(a) for compounds of formula (I) where Y is C=O, reaction of a compound of formula (II):



(II)

30

in which R¹, X, m, n, p and q are as defined in formula (I) or are protected derivatives thereof, with a compound of formula (III):



in which Ar, R³, R⁴ and r are as defined in formula (I) or are protected derivatives thereof,
or

- 5 (b) for compounds of formula (I) where Y is SO₂, reaction of a compound of formula (II) with a compound of formula (IV):



- 10 in which Ar, R³, R⁴ and r are as defined in formula (I) or are protected derivatives thereof,
or

(c) for compounds of formula (I) where Y is CONH, reaction of a compound of formula (II) with a carbonyl source such as phosgene or triphosgene and an amine

- 15 Ar-(CR³R⁴)_r-NH₂ or by treating with an isocyanate Ar-(CR³R⁴)_r-NCO, or

(d) for compounds of formula (I) where r is 0 and Y is CH₂, reaction of a compound of formula (II) with a compound ArCHO by reductive amination,

- 20 and optionally thereafter,

- removing any protecting groups
- forming a pharmaceutically acceptable salt.

- 25 The reaction between compounds (II) and (III) may be carried out using standard coupling conditions for example using peptide coupling reagents such as HOBt, DCC PyBrop, or via an acid chloride in the presence of a base such as triethylamine in an inert solvent.


Reaction of compounds (II) and (IV) can be carried out in the presence of a base such as triethylamine or pyridine in an aprotic solvent such as dichloromethane.

30 Process (d) can be carried out by reductive amination using reagents such as solid supported cyanoborohydride resin, catalytic acetic acid in aprotic solvent such as dichloromethane or NMP, or alternatively sodium triacetoxyborohydride in dichloromethane with catalytic acetic acid.

35


PRV 03-07-21

Compounds of formula (II) where X is NH₂ or SH may be prepared from compounds of formula (V)



(V)


where R¹ and m are as defined above and L is a leaving group by reaction with a compound of formula (VI):



(VI)

where X is NH₂ or SH, PG is a protecting group, L is a leaving group, and n, p and q are as defined in formula (I), and where X is SH optionally oxidising the resulting compound of formula (II). The reaction can be carried out using an aprotic base such as triethylamine or Hunig's base in a suitable solvent such as dichloromethane. Suitable protecting groups PG include acid labile groups such as tBoc. Compounds of formula (II) where X is S can be oxidised using oxone or mCPBA under controlled conditions to give the corresponding compounds where X is SO or SO₂.

Compounds of formula (II) where X is O can be prepared by reacting a compound of formula (V) as defined above with a compound of formula (VII):



(VII)

in which PG is a protecting group, L is a leaving group, and n, p and q and are as defined in formula (I). Preferably L in compound (V) is halide or a triflate, the reaction being carried out in the presence of a base such as sodium hydride or potassium t-butoxide. The group PG is an acid labile group such as t-Boc.

It will be appreciated by those skilled in the art that in the processes of the present invention certain functional groups such as hydroxyl or amino groups in the starting reagents or intermediate compound may need to be protected by protecting groups. Thus, the preparation of the compound of formula (I) may involve, at an appropriate stage, the removal of one or more protecting groups. The protection and deprotection of functional groups is fully described in 'Protective Groups in Organic Chemistry', edited by J. W. F. McOmie, Plenum Press (1973), and 'Protective Groups in Organic Synthesis', 2nd edition, T. W. Greene & P. G. M. Wuts, Wiley-Interscience (1991).

The compounds of formula (I) above may be converted to a pharmaceutically acceptable salt or solvate thereof, preferably a basic addition salt such as sodium, potassium, calcium, aluminium, lithium, magnesium, zinc, benzathine, chlorprocaine, choline, diethanolamine, ethanolamine, ethyldiamine, meglumine, tromethamine or procaine, or an acid addition salt such as a hydrochloride, hydrobromide, phosphate, acetate, fumarate, maleate, tartrate, citrate, oxalate, methanesulphonate or *p*-toluenesulphonate.

The compounds of formula (I) have activity as pharmaceuticals, in particular as modulators of histamine H₄, and may be used in the treatment (therapeutic or prophylactic) of conditions/diseases in human and non-human animals including:

- (1) (the respiratory tract) obstructive airways diseases including chronic obstructive pulmonary disease (COPD); asthma, such as bronchial, allergic, intrinsic, extrinsic and dust asthma, particularly chronic or inveterate asthma (e.g. late asthma and airways hyper-responsiveness); bronchitis; acute, allergic, atrophic rhinitis and chronic rhinitis including rhinitis caseosa, hypertrophic rhinitis, rhinitis purulenta, rhinitis sicca and rhinitis medicamentosa; membranous rhinitis including croupous, fibrinous and pseudomembranous rhinitis and scrofulous rhinitis; seasonal rhinitis including rhinitis nervosa (hay fever) and vasomotor rhinitis; sarcoidosis, farmer's lung and related diseases, fibroid lung and idiopathic interstitial pneumonia;

- (2) **(bone and joints)** gout, rheumatoid arthritis, seronegative spondyloarthropathies (including ankylosing spondylitis, psoriatic arthritis and Reiter's disease), Behcet's disease, Sjogren's syndrome and systemic sclerosis;
- (3) **(skin)** pruritis, scleroderma, otitis, psoriasis, atopic dermatitis, contact dermatitis and other eczematous dermatides, seborrhoetic dermatitis, Lichen planus, Pemphigus, bullous Pemphigus, Epidermolysis bullosa, urticaria, angiodermas, vasculitides, erythemas, cutaneous eosinophilias, uveitis, Alopecia areata and vernal conjunctivitis, lupus;
- (4) **(gastrointestinal tract)** Coeliac disease, proctitis, eosinophilic gastro-enteritis, mastocytosis, inflammatory bowel diseases such as Crohn's disease, ulcerative colitis, ileitis and enteritis, food-related allergies which have effects remote from the gut, e.g., migraine, rhinitis and eczema;
- (5) **(central and peripheral nervous system)** Neurodegenerative diseases and dementia disorders, e.g. Alzheimer's disease, amyotrophic lateral sclerosis and other motor neuron diseases, Creutzfeldt-Jacob's disease and other prion diseases, HIV encephalopathy (AIDS dementia complex), Huntington's disease, frontotemporal dementia, Lewy body dementia and vascular dementia; polyneuropathies, e.g. Guillain-Barré syndrome, chronic inflammatory demyelinating polyradiculoneuropathy, multifocal motor neuropathy, plexopathies; CNS demyelination, e.g. multiple sclerosis, acute disseminated/haemorrhagic encephalomyelitis, and subacute sclerosing panencephalitis; neuromuscular disorders, e.g. myasthenia gravis and Lambert-Eaton syndrome; spinal disorders, e.g. tropical spastic paraparesis, and stiff-man syndrome; paraneoplastic syndromes, e.g. cerebellar degeneration and encephalomyelitis; CNS trauma; migraine; stroke and connective tissue diseases such as meningitis
- (6) **(other tissues and systemic disease)** hepatitis, vasculitis, spondyloarthropathies, vaginitis, glomerulonephritis, myositis, atherosclerosis, Acquired Immunodeficiency Syndrome (AIDS), lupus erythematosus, systemic lupus, erythematosus, Hashimoto's thyroiditis, type I diabetes, nephrotic syndrome, eosinophilia fasciitis, hyper IgE syndrome, lepromatous

leprosy, and idiopathic thrombocytopenia pupura; post-operative adhesions, and sepsis.

5 (7) (allograft and xenograft rejection) acute and chronic following, for example, transplantation of kidney, heart, liver, lung, bone marrow, skin and cornea; and chronic graft versus host disease;

10 (8) Cancer, carcinoma & tumour metastasis, including that of the bladder, breast, colon, kidney, liver, lung, ovary, pancreas, stomach, cervix, thyroid and skin, especially non-small cell lung cancer (NSCLC), malignant melanoma, prostate cancer and squamous sarcoma. Hematopoietic tumors of lymphoid lineage, including acute lymphocytic leukemia, B cell lymphoma and Burketts lymphoma, Hodgkins Lymphoma, Acute Lymphoblastic Leukemia. Hematopoietic tumors of myeloid lineage, including acute and chronic
15 myelogenous leukemias and promyelocytic leukemia. Tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma, and other tumors, including melanoma, seminoma, tetratocarcinoma, neuroblastoma and glioma.

20 (9) All diseases that result from a general imbalance of the immune system and resulting in increased atopic inflammatory reactions.

(10) Cystic fibrosis, re-perfusion injury in the heart, brain, peripheral limbs and other organs.

25 (11) Burn wounds & chronic skin ulcers

(12) Reproductive Diseases (e.g. Disorders of ovulation, menstruation and implantation, Pre-term labour, Endometriosis)

30 (13) thrombosis

(14) infectious diseases such as HIV infection and other viral infections, bacterial infections.

35 Thus, the present invention provides a compound of formula (IA), or a pharmaceutically-acceptable salt or solvate thereof, as hereinbefore defined for use in therapy.

Preferably the compounds of the invention are used to treat respiratory diseases. It is preferred that the compound of the invention is used to treat asthma and rhinitis, especially asthma.

5

In a further aspect, the present invention provides the use of a compound of formula (IA), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for use in therapy.

10

In a still further aspect, the present invention provides the use of a compound of formula (IA), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for the treatment of human diseases or conditions in which modulation of histamine H4 receptor activity is beneficial.

15

In the context of the present specification, the term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The terms "therapeutic" and "therapeutically" should be construed accordingly.

20

The invention still further provides a method of treating a histamine H4 mediated disease, which comprises administering to a patient a therapeutically effective amount of a compound of formula (I) or (IA), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined.

25

The invention also provides a method of treating a respiratory disease, such as asthma and rhinitis, especially asthma, in a patient suffering from, or at risk of, said disease, which comprises administering to the patient a therapeutically effective amount of a compound of formula (I) or (IA), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined.

30

For the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disorder indicated.

35

The compound of formula (I) and pharmaceutically acceptable salts and solvates thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in which the formula (I) compound/salt/solvate (active

ingredient) is in association with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of administration, the pharmaceutical composition will preferably comprise from 0.05 to 99 %w (per cent by weight), more preferably from 0.05 to 80 %w, still more preferably from 0.10 to 70 %w, and even more preferably from 0.10 to 50 %w, of active ingredient, all percentages by weight being based on total composition.


The present invention also provides a pharmaceutical composition comprising a compound of formula (IA), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined, in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

The invention further provides a process for the preparation of a pharmaceutical composition of the invention which comprises mixing a compound of formula (IA), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined, with a pharmaceutically acceptable adjuvant, diluent or carrier.

The compounds of the invention can be administered in combination with other agents such as long-acting β -agonists.


The pharmaceutical compositions may be administered topically (e.g. to the lung and/or airways or to the skin) in the form of solutions, suspensions, heptafluoroalkane aerosols and dry powder formulations; or systemically, e.g. by oral administration in the form of tablets, capsules, syrups, powders or granules, or by parenteral administration in the form of solutions or suspensions, or by subcutaneous administration or by rectal administration in the form of suppositories or transdermally. Preferably the compound of the invention is administered orally.

The following examples illustrate the invention.

PREPARATION OF INTERMEDIATES**1-Piperidinecarboxylic acid, 4-[[1-(triphenylmethyl)-1H-imidazol-4-yl]methoxy]-, 1,1-dimethylethyl ester**


To a solution of 4-hydroxy-1-piperidinecarboxylic acid, 1,1-dimethylethyl ester (9.5g, 0.047mol) in dry N-methylpyrrolidine (NMP) (50ml) was added sodium hydride (60% in oil) portionwise (1.88g, 0.047mol). The mixture was allowed to stir for 30 minutes, then 4-(chloromethyl)-1-(triphenylmethyl)-1H-imidazole (ref : WO0244141) (16.8g, 0.047mol) was added and the mixture heated to 50°C for 30 min, poured into ice water and extracted with ethyl acetate. The organic extracts were washed with water and purified by flash column chromatography eluting with 1% methanolic ammonia/dichloromethane to give a solid (5.5g). 300 MHz ¹H NMR (CDCl₃) 7.41 (1H, d), 7.32-7.13 (15H, m), 6.81 (1H, bs), 4.49 (2H, s), 3.82-3.77 (2H, m), 3.62-3.54 (1H, m), 3.05-2.96 (2H, m), 1.85-1.76 (2H, m), 1.57-1.48 (2H, m), 1.45 (9H, s)

20 1-Piperidinecarboxylic acid, 4-[[5-methyl-1-(triphenylmethyl)-1H-imidazol-4-yl]methoxy]-, 1,1-dimethylethyl ester




This was prepared by the method of Example 1 from 4-(chloromethyl)-5-methyl-1-(triphenylmethyl)-1H-imidazole (ref: European Journal of Medicinal Chemistry (1990), 25(7), 557). 300 MHz ¹H NMR (CDCl₃) 7.34-7.29 (15H, m), 7.26 (1H, d), 4.47 (2H, s), 3.82-3.77 (2H, m), 3.62-3.54 (1H, m), 3.05-2.96 (2H, m), 2.25 (3H, s), 1.85-1.76 (2H, m), 1.57-1.48 (2H, m), 1.45 (9H, s)

4-[[1-(Triphenylmethyl)-1H-imidazol-4-yl]methoxy]-piperidine



Trimethylsilyliodide (3.3ml, 1 eq.) was added to a solution of 1-piperidinecarboxylic acid, 4-[[1-(triphenylmethyl)-1H-imidazol-4-yl]methoxy]-, 1,1-dimethylethyl ester (12g, 0.023mol) at 0 – 5°C, stirred at this temperature for 30minutes, quenched with ice cold sodium bicarbonate solution and the organic layer separated, dried over sodium sulphate and evaporated. The residue was purified by flash column chromatography eluting with 5% methanolic ammonia/dichloromethane to give a solid (6.9g). 300 MHz ¹H NMR (CDCl₃) 7.41 (1H, d), 7.11-7.39 (15H, m), 6.81 (1H, d), 4.48 (2H, s), 3.53 (1H, m), 3.41 (2H, s), 3.08 (2H, m), 2.67 (2H, m), 1.99 (2H, m), 1.51 (2H, m)

4-[[5-Methyl-1-(triphenylmethyl)-1H-imidazol-4-yl]methoxy]-piperidine



This was prepared by the method of example 2. MS (+APCI) m/z 438 ($M+H^+$)


5

PREPARATION OF FINAL PRODUCTS

Example 1

10

4-(1*H*-Imidazol-4-ylmethoxy)-1-(1-oxo-3-phenylbutyl)-piperidine




- 15 **Bromo-tris-pyrrolidinophosphonium hexafluorophosphate (PyBrop™)** (0.55g, 1.18mmol) was added to a solution of 4-[[1-(triphenylmethyl)-1*H*-imidazol-4-yl]methoxy]-piperidine (0.5g, 1.18mmol) and 3-phenylbutanoic acid (0.24g, 1.18mmol), Hunig's base (1 ml) in dry NMP. The mixture was stirred at room temp for 16h, evaporated and the residue dissolved in methanol, filtered through sulphonic acid ion exchange resin eluting with
- 20 methanol followed by methanolic ammonia, and evaporated. The residue was purified by reverse phase HPLC on an Xterra™ column eluting with acetonitrile/aqueous ammonium hydroxide to give the product as a white solid (0.052g). MS (+APCI) m/z 328 ($M+H^+$). 400 MHz ¹H NMR (d₆-DMSO) (@125C – spectrum at room temperature complicated due to rotamers) 7.54 (1H, s), 7.29-7.24 (3H, m), 7.19-7.14 (2H, m), 6.91 (1H, s), 4.42 (2H, s).

3.75-3.58 (2H, m), 3.30-3.08 (2H, m), 2.8 (1H, br.s), 2.61 (2H, dd), 2.53 (4H dd), 1.80-1.70 (2H, m), 1.47-1.33 (H, m), 1.25 (2H, d)

Example 2


4-(1*H*-Imidazol-4-ylmethoxy)-1-[[4-(trifluoromethyl)phenyl]acetyl]-piperidine



The title compound was prepared using the method of Example 1 with 4-(trifluoromethyl)phenylacetic acid : MS (+APCI) m/z 368 ($M+H^+$). 400 MHz 1H NMR (d_6 -DMSO) (@ 125C – spectrum at room temperature complicated due to rotamers) 7.63 (2H, d), 7.57 (1H, s), 7.44 (2H, d), 7.02 (1H, br.s), 4.44 (2H, s), 3.83 (2H, s), 3.80-3.73 (1H, m), 3.70-3.62 (2H, m), 3.31-3.22 (2H, m), 2.88 (2H, br.s), 1.83-1.73 (2H, br.m), 1.50-1.41 (2H, br.m)

Example 3






1-[2-(4-Hydroxyphenyl)-1-oxopropyl]-4-[(5-methyl-1*H*-imidazol-4-yl)methoxy]-piperidine













This was prepared by the method of Example 1 using 2-(4-hydroxyphenyl)propanoic acid and 4-[[5-methyl-1-(triphenylmethyl)-1*H*-imidazol-4-yl]methoxy]-piperidine. MS (+APCI) m/z 344 ($M+H^+$). 400 MHz 1H NMR (d_6 -DMSO) 11.45 (1*H*, s), 8.96 (1*H*, s), 7.32 (1*H*, s), 6.96 (2*H*, m), 6.68 (2*H*, m), 4.30 (2*H*, s), 3.93 (1*H*, s), 3.71 (2*H*, m), 3.46 (1*H*, m), 3.25 (2*H*, m), 2.09 (3*H*, s), 1.63 (2*H*, m), 1.23 (5*H*, m)

The compounds in Table 1 were prepared using the method of Example 1 with the appropriate acid :

Table 1

Example structure	Example no.	Name	$M+H^+$ (+APCI)
	4	piperidine, 1-[(4-fluorophenyl)acetyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	317
	5	piperidine, 1-[(2-chlorophenyl)acetyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	333
	6	piperidine, 1-[(4-chlorophenyl)acetyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	333
	7	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-(phenylacetyl)-	299
	8	piperidine, 1-(4-cyclohexylbenzoyl)-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	367


	9	piperidine, 1-[(3,4-dichlorophenyl)acetyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	367
	10	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-[(4-methylphenyl)acetyl]-	311
	11	piperidine, 1-[(3,4-difluorophenyl)acetyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	327
	12	piperidine, 1-[(2,4-difluorophenyl)acetyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	317
	13	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-[(4'-propyl[1,1'-biphenyl]-4-yl)carbonyl]-	355
	14	piperidine, 1-[2-(4-hydroxyphenyl)-1-oxopropyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	329
	15	piperidine, 1-[(2 <i>E</i>)-3-(3,4-dichlorophenyl)-1-oxo-2-propenyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	379
	16	piperidine, 1-[3-(2,4-dichlorophenyl)-1-oxopropyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	381
	17	piperidine, 1-[(2,4-dichlorophenyl)acetyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	367

5 **Example 18****1-[(2-Bromophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine**

10

To a solution of 4-[[1-(triphenylmethyl)-1*H*-imidazol-4-yl]methoxy]-piperidine (0.5g, 1.18mmol) and 2-bromobenzaldehyde (0.218g, 1.18mmol) in 10% acetic acid/*N*-methylpyrrolidine (10ml) was added (polystyrylmethyl)trimethylammonium cyanoborohydride resin (4.2mmol/g equivalent, 300mg). The mixture was stirred 16h at room temperature, the resin removed by filtration and the solution evaporated. The residue was purified by reserve phase HPLC on an Xterra™ column using aqueous ammonium acetate/acetonitrile as eluant to give the title compound (0.058g). MS (+APCI) *m/z* 350 (*M*+*H*⁺). 400 MHz ¹H NMR (d₆-DMSO) 7.59-7.56 (2H, m), 7.46 (1H, d), 7.36 (1H, t), 7.19 (1H, t), 7.05-6.8 (2H, br.m), 4.37 (2H, br.s), 3.50 (2H, s), 3.40 (1H, br.s), 2.71-2.66 (2H, m), 2.18-2.12 (2H, m), 1.84-1.81 (2H, m), 1.50-1.41 (2H, m)









15
20**Example 19****1-[(3-Bromo-2-thienyl)methyl]-4-[(5-methyl-1*H*-imidazol-4-yl)methoxy]- piperidine**





















- This was prepared by the method of Example 18 using 3-bromothiophene-2-carboxaldehyde and 4-[[5-methyl-1-(triphenylmethyl)-1*H*-imidazol-4-yl]methoxy]piperidine. MS (+APCI) m/z 372 ($M+H^+$). 400 MHz 1H NMR (d_6 -DMSO) 7.57 (1H, d, J 5.4 Hz), 7.40 (1H, s), 7.02 (1H, d, J 5.4 Hz), 4.32 (2H, s), 3.61 (2H, s), 3.29 (1H, m), 2.73 (2H, m), 2.17 (2H, m), 2.11 (3H, s), 1.82 (2H, m), 1.42 (2H, m)
- 10 The compounds in Table 2 were prepared using the method of Example 18 with the appropriate acid :










Table 2










Example structure	Example no.	Name	$M+H^+$ (+APCI)
	20	piperidine, 1-[(3-bromo-2-thienyl)methyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	355
	21	piperidine, 1-[(4-ethynylphenyl)methyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	295










	22	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-[[3-(4-methylphenoxy)phenyl]methyl]-	377
	23	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-[[4-(2-propenyloxy)phenyl]methyl]-	327
	24	phenol, 4-[[4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-	287
	25	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-[(2-methoxyphenyl)methyl]-	301
	26	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-[[3-(4-methoxyphenoxy)phenyl]methyl]-	393
	27	piperidine, 1-[(2,3-dichlorophenyl)methyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	339
	28	piperidine, 1-[(2-chloro-4-fluorophenyl)methyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	323
	29	piperidine, 1-(2-dibenzofuranylmethyl)-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	361










	30	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-[[2-(methylthio)phenyl]methyl]-	317
	31	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-(thieno[2,3- <i>b</i>][1]benzothien-2-ylmethyl)-	383
	32	piperidine, 1-[(2-chloro-5-nitrophenyl)methyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	350
	33	1 <i>H</i> -pyrrole, 2-[[4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-1-[(4-methylphenyl)sulfonyl]-	414
	34	phenol, 2-ethoxy-6-[[4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-	331
	35	piperidine, 1-(1,3-benzodioxol-5-ylmethyl)-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	315
	36	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-[[4-(phenylmethoxy)phenyl]methyl]-	377
	37	piperidine, 1-[[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	357










	38	piperidine, 1-[(4-bromophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	349
	39	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[(4-methylphenyl)methyl]-	285
	40	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-(2-thienylmethyl)-	277
	41	piperidine, 1-[(4-chlorophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	305
	42	piperidine, 1-[(2-chloro-6-fluorophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	323
	43	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[(3-methyl-2-thienyl)methyl]-	291
	44	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-(2-naphthalenylmethyl)-	321
	45	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-(1-naphthalenylmethyl)-	321
	46	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[(2-nitrophenyl)methyl]-	316
	47	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-(3-thienylmethyl)-	277










	48	piperidine, 1-([1,1'-biphenyl]-4-ylmethyl)-4-(1H-imidazol-4-ylmethoxy)-	347
	49	piperidine, 1-[(2,5-difluorophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	307
	50	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[(3-phenoxyphenyl)methyl]-	363
	51	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[(3-methylphenyl)methyl]-	285
	52	piperidine, 1-(2-furanylmethyl)-4-(1H-imidazol-4-ylmethoxy)-	261
	53	piperidine, 1-[(2,6-dichlorophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	339
	54	piperidine, 1-[(4-fluorophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	289
	55	piperidine, 1-[(3-fluorophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	289
	56	piperidine, 1-(3-furanylmethyl)-4-(1H-imidazol-4-ylmethoxy)-	261




	57	piperidine, 1-[(4-ethylphenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	299
	58	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[(2-methylphenyl)methyl]-	285
	59	piperidine, 1-[(3-chlorophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	305
	60	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[(5-methyl-2-thienyl)methyl]-	291
	61	piperidine, 1-[(4-bromo-2-thienyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	355
	62	piperidine, 1-[(2,2'-bithiophen]-5-ylmethyl)-4-(1H-imidazol-4-ylmethoxy)-	359
	63	phenol, 3,5-dichloro-2-[[4-(1H-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-	355
	64	piperidine, 1-[(3,4-difluorophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	307
	65	piperidine, 1-[(3,5-difluorophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	307

	66	piperidine, 1-[(6-chloro-1,3-benzodioxol-5-yl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	349
	67	piperidine, 1-[[4-[4-(1,1-dimethylethyl)-2-thiazolyl]phenyl]methyl]-4-(1H-imidazol-4-ylmethoxy)-	410
	68	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[(1-methyl-1H-pyrrol-2-yl)methyl]-	274
	69	1H-indole, 3-[[4-(1H-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-1-(phenylmethyl)-	400
	70	piperidine, 1-[(5-chloro-2-thienyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	311
	71	piperidine, 1-(1,3-benzodioxol-4-ylmethyl)-4-(1H-imidazol-4-ylmethoxy)-	315
	72	2-thiophenecarbonitrile, 3-[[4-[[4-(1H-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]phenoxy]methyl]-	408
	73	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[[5-(phenylethynyl)-2-thienyl]methyl]-	377
	74	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[[5-(4-nitrophenyl)-2-thienyl]methyl]-	382

	75	furanyl]methyl]- piperidine, 4-(1 <i>H</i> - imidazol-4-ylmethoxy)-1- [[5-(3-nitrophenyl)-2- furanyl]methyl]-	382
	76	piperidine, 1-[(4-chloro- 1 <i>H</i> -pyrazol-3-yl)methyl]- 4-(1 <i>H</i> -imidazol-4- ylmethoxy)-	295
	77	piperidine, 1-[(4-bromo- 1-methyl-1 <i>H</i> -pyrazol-3- yl)methyl]-4-(1 <i>H</i> - imidazol-4-ylmethoxy)-	353
	78	piperidine, 1-[(4-bromo- 1 <i>H</i> -pyrazol-3-yl)methyl]- 4-(1 <i>H</i> -imidazol-4- ylmethoxy)-	339
	79	benzonitrile, 2-[[4-(1 <i>H</i> - imidazol-4-ylmethoxy)-1- piperidinyl]methyl]-	296
	80	piperidine, 4-(1 <i>H</i> - imidazol-4-ylmethoxy)-1- [(4-iodophenyl)methyl]-	397
	81	piperidine, 1-[(5-ethyl-2- thienyl)methyl]-4-(1 <i>H</i> - imidazol-4-ylmethoxy)-	305
	82	piperidine, 4-(1 <i>H</i> - imidazol-4-ylmethoxy)-1- [[5-(methylthio)-2- thienyl]methyl]-	323
	83	piperidine, 1-[[1-(3,5- dichlorophenyl)-1 <i>H</i> - pyrrol-2-yl]methyl]-4- (1 <i>H</i> -imidazol-4- ylmethoxy)-	404

	84	piperidine, 1-[[1-(4-chlorophenyl)-1H-pyrrol-2-yl]methyl]-4-(1H-imidazol-4-ylmethoxy)-	370
	85	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[[4-(phenylethynyl)-2-thienyl]methyl]-	377
	86	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[(3-phenoxy-2-thienyl)methyl]-	369
	87	piperidine, 1-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-4-(1H-imidazol-4-ylmethoxy)-	373
	88	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[(4-propoxyphenyl)methyl]-	329
	89	phenol, 2-[[4-(1H-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-	287
	90	piperidine, 1-[(2,4-difluorophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	307
	91	2-thiophenecarbonitrile, 3-[[4-(1H-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-	302
	92	piperidine, 1-(benzo[b]thien-3-ylmethyl)-4-(1H-imidazol-4-ylmethoxy)-	327

	93	pyridine, 2-chloro-3-[[4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-	306
	94	phenol, 3-[[4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-2-(2-propenyl)-	327
	95	piperidine, 1-[(4-chloro-3-fluorophenyl)methyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	323
	96	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-[[4-(trifluoromethoxy)phenyl]methyl]-	355
	97	piperidine, 1-[(2,6-difluorophenyl)methyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	307
	98	piperidine, 1-[(4-bromo-2-fluorophenyl)methyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	367
	99	piperidine, 1-[(2,2-difluoro-1,3-benzodioxol-5-yl)methyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	351
	100	piperidine, 1-[(4-butoxyphenyl)methyl]-4-(1 <i>H</i> -imidazol-4-ylmethoxy)-	343
	101	piperidine, 4-(1 <i>H</i> -imidazol-4-ylmethoxy)-1-[(2,3,5-	373

		trichlorophenyl)methyl]-	
	102	piperidine, 1-[(2,5-dichlorophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	339
	103	piperidine, 4-(1H-imidazol-4-ylmethoxy)-1-[[2-(trifluoromethyl)phenyl]methyl]-	339
	104	piperidine, 1-[(4-chloro-2-nitrophenyl)methyl]-4-(1H-imidazol-4-ylmethoxy)-	350

Pharmacological Data

H4-CHO FLUOROMETRIC IMAGING PLATE READER (FLIPR) ASSAY

5 FLIPR was employed to measure the intracellular calcium mobilisation to H4 receptor activation by histamine. CHO-K1 cells expressing the human recombinant H4 receptor with G α 16 were purchased from Euroscreen and used in the experiments to identify H4 antagonists. The same protocol was used with the human H3-CHO cell line (Euroscreen) to determine selectivity of the H4 antagonists.

10

Briefly, the FLIPR protocol detects changes in $[Ca^{2+}]_i$ using Fluo-3AM loaded cells (Schroeder & Neagle. FLIPR: A new instrument for accurate, high throughput optical screening. *J. Biomol. Screening*: 1(2), 75-80, 1996.). The H4-CHO cells were cultured routinely in T225cm² tissue culture flasks as monolayers in NUT Hams (with 1% (v/v) 15 Glutamine) supplemented with 10% (v/v) heat inactivated foetal bovine serum and grown under Geneticin (1mg/ml) antibiotic selection & 1mg/ml Zeocin selection. Cultures were maintained at 37°C in a humidified atmosphere of 5% CO₂ and passaged every 3 days.


H4-CHO cells were seeded at 10,000 cells/well (384 FLIPR plate) 18-24hr before the 20 experiment. Cells were washed to remove medium and replaced with loading buffer for 1.5 hrs. The loading buffer contains Hanks balance salt solution (Sigma), HEPES (20 mM), probenecid (2.5 mM) and Fluo 3-AM (4 μ M) /Brilliant Black at pH7.4. The EC50 of histamine was determined on the day of the experiment and 2X EC50 was chosen as the dose to test compounds against. ATP stimulation was included in the FLIPR assay to 25 exclude any non-selective antagonists.

Step by step guide to FLIPR assay

1. Cells were harvested using 1x dissociation solution and plated onto poly-D-lysine coated FLIPR 384 plates at 1.0×10^4 cells per well 18-24 hours prior to experiment.

2. Media was removed from the cells by tipping the plates and gently blotted onto tissue to remove any excess medium.
3. 30 μ l loading buffer was added to all wells and plates were incubated for 90 min at 37°C.
- 5 4. 96 well histamine EC₅₀ plate was made and then 40 μ l was indexed into 4 quadrants in a 384 well plate.
5. 96 well compound vehicle (1% DMSO) plate was made and indexed into a quadrant of a 384 well plate.
6. Plates transferred to FLIPR and run using protocol attached (see Appendix 1)
- 10 7. EC₅₀ for histamine was calculated.
8. 96 well histamine plate (x10 EC₅₀) was made and then 60 μ l was indexed into 4 quadrants in a 384 well plate.
9. Each 96 well compound plate was made and indexed into a quadrant of a 384 well plate.
- 15 10. ATP plate was made in a 96 well plate and then 60 μ l was indexed into 4 quadrants in a 384 well plate.
11. Plates transferred to FLIPR and run using protocol attached (see Appendix 1)
 - Cell media (but not cells) removed from FLIPR 384 plate
 - 30 μ l of loading buffer added to FLIPR 384 plate
 - 20 • 10 μ l compound added to cell plate
 - Reads taken for 5min to determine compound effects
 - 10 μ l histamine added to cell plate
 - Reads taken to determine histamine response
 - 10 μ l ATP added to cell plate
 - 25 • Reads taken for 5min to determine ATP response
12. Final assay concentrations
 - Compound concentration range = 30 μ M to 0.01 μ M
 - histamine= 2x calculated EC₅₀
 - ATP = 11 μ M

5

Example of FLIPR traceA = Compound additionB = Histamine additionC = ATP addition


10

Appendix 1**384 well FLIPR protocol****General**

5	Exposure	0.4
	Filter	1
	Presoak	None
	Second Sequence	Yes
	Third Sequence	YES
10	Auto prompt for notes	No
	Auto print	Yes
	Auto export time sequence	Yes
	Auto export stats	Yes
First sequence (Compound)		
15	Initial interval	20 readings every 2sec
	Second interval	27 readings every 10sec
	Fluid Addition	
	Addition Active	Yes
20	Volume	10.0
	After Sample	5
	Height	25
	Speed	15.0
	Mix	No
Second sequence (histamine)		
25	Initial interval	20 readings every 2sec
	Second interval	12 readings every 10sec
	Fluid Addition	
	Addition Active	Yes
30	Volume	10.0
	After Sample	3
	Height	35
	Speed	15.0
	Mix	No
Third sequence (ATP)		
35	Initial interval	20 readings every 2sec
	Second interval	12 readings every 10sec
	Fluid Addition	
	Addition Active	Yes
40	Volume	10.0
	After Sample	3
	Height	35
	Speed	40.0
	Mix	No
Pipetting		
45	Mix Volume	0.0
	Mix Cycles	0
	Leave tips in well	No
	Remove fluid after addition	No
	Stage heated to 35°C	
50	The compounds of the examples have an IC ₅₀ vs H4 of <10 micromolar.	

CLAIMS

1. A compound of formula (I) and pharmaceutically acceptable salts and solvates thereof
 5 for use in the manufacture of a medicament for the treatment of diseases mediated by histamine H3 and H4:



(I)

in which:

Ar is an aryl group, a 5-7 membered heteraromatic ring containing 1-4 heteroatoms
 15 selected from nitrogen, oxygen or sulphur, or a bicyclic or tricyclic heteraromatic ring containing 1-4 heteroatoms selected from nitrogen, oxygen or sulphur, each of which can be optionally substituted by 1-3 groups selected from C₁₋₆ alkyl, C₁₋₆ alkylthio, C₁₋₆ alkoxy, halogen, cyano, CF₃, OCF₃, C₃₋₆ cycloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₂₋₆ alkenyloxy, hydroxyl, nitro, tosyl, thienyl, benzyl, phenyl, nitrophenyl,

20 R¹ is hydrogen or C₁₋₆ alkyl;

X is O, NR², CH₂ or SO_x

R² is C₁₋₆ alkyl;

25 x is 0, 1 or 2;

Y is CH₂, C=O, SO₂, or (C=O)NH;

30 Z is (CR³R⁴)_r, or Y and Z together form a CH=CH group;

m and n are independently 0, 1, 2 or 3;

p and q are independently 0, 1 or 2;

r is 0, 1, 2, 3, or 4 and

5 R^3 and R^4 are independently hydrogen or C_{1-6} alkyl.

2. A compound according to claim 1 in which Ar^1 is phenyl optionally substituted as defined above. Preferred substituents include halogen such as iodo, chloro and fluoro, cyclohexyl, methyl, ethyl, propyl, t-butyl, ethynyl, propenyloxy, hydroxyl, methoxy, nitro,
10 tosyl, trifluoromethyl, thienyl, benzyl, cyano, phenylethynyl, nitrophenyl, methylthio, propoxy, butoxy, 2-propenyl, or trifluomethoxy.

3. A compound according to claim 1 or 2 in which R^1 is hydrogen or methyl.

15 4. A compound according to any one of claims 1 to 3 in which X is O.

5. A compound according to any one of claims 1 to 4 in which Y is CH_2 or $C=O$ and Z is CH_2 , $CHMe$, CH_2CHMe or Y and Z form a $CH=CH$ group.

20 6. A compound according to any one of claims 1 to 5 in which m is 1 and n is 0.

7. A compound according to any one of claims 1 to 6 in which p and q are both 1.

8. A compound of formula (I) selected from:

25 4-(1*H*-Imidazol-4-ylmethoxy)-1-(1-oxo-3-phenylbutyl)-piperidine

4-(1*H*-Imidazol-4-ylmethoxy)-1-[[4-(trifluoromethyl)phenyl]acetyl]-piperidine

1-[2-(4-Hydroxyphenyl)-1-oxopropyl]-4-[(5-methyl-1*H*-imidazol-4-yl)methoxy]-
piperidine

1-[(4-fluorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine

30 1-[(2-chlorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine

1-[(4-chlorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine

4-(1*H*-imidazol-4-ylmethoxy)-1-(phenylacetyl)-piperidine

1-(4-cyclohexylbenzoyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine

1-[(3,4-dichlorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine


- 4-(1*H*-imidazol-4-ylmethoxy)-1-[(4-methylphenyl)acetyl]-piperidine
 1-[(3,4-difluorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-[(2,4-difluorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 4-(1*H*-imidazol-4-ylmethoxy)-1-[(4'-propyl[1,1'-biphenyl]-4-yl)carbonyl]-piperidine
 5 1-[2-(4-hydroxyphenyl)-1-oxopropyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-[(2*E*)-3-(3,4-dichlorophenyl)-1-oxo-2-propenyl]-4-(1*H*-imidazol-4-ylmethoxy)-
 piperidine
 1-[3-(2,4-dichlorophenyl)-1-oxopropyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-[(2,4-dichlorophenyl)acetyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 10 1-[(2-Bromophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-[(3-Bromo-2-thienyl)methyl]-4-[(5-methyl-1*H*-imidazol-4-yl)methoxy]-piperidine
 1-[(3-bromo-2-thienyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-[(4-ethynylphenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 4-(1*H*-imidazol-4-ylmethoxy)-1-[[3-(4-methylphenoxy)phenyl]methyl]-piperidine
 15 4-(1*H*-imidazol-4-ylmethoxy)-1-[[4-(2-propenyloxy)phenyl]methyl]-piperidine
 4-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-phenol
 4-(1*H*-imidazol-4-ylmethoxy)-1-[(2-methoxyphenyl)methyl]-piperidine
 4-(1*H*-imidazol-4-ylmethoxy)-1-[[3-(4-methoxyphenoxy)phenyl]methyl]-piperidine
 1-[(2,3-dichlorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 20 1-[(2-chloro-4-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-(2-dibenzofuranylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 4-(1*H*-imidazol-4-ylmethoxy)-1-[[2-(methylthio)phenyl]methyl]-piperidine
 4-(1*H*-imidazol-4-ylmethoxy)-1-(thieno[2,3-*b*][1]benzothien-2-ylmethyl)-piperidine
 1-[(2-chloro-5-nitrophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 25 1*H*-pyrrole, 2-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-1-[(4-
 methylphenyl)sulfonyl]-
 2-ethoxy-6-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-phenol
 1-(1,3-benzodioxol-5-ylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 4-(1*H*-imidazol-4-ylmethoxy)-1-[[4-(phenylmethoxy)phenyl]methyl]-piperidine
 30 1-[[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine

- 1-[(4-bromophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(4-methylphenyl)methyl]-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-(2-thienylmethyl)-piperidine
1-[(4-chlorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
5 1-[(2-chloro-6-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(3-methyl-2-thienyl)methyl]-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-(2-naphthalenylmethyl)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-(1-naphthalenylmethyl)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(2-nitrophenyl)methyl]-piperidine
10 4-(1*H*-imidazol-4-ylmethoxy)-1-(3-thienylmethyl)-piperidine
1-([1,1'-biphenyl]-4-ylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(2,5-difluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(3-phenoxyphenyl)methyl]-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(3-methylphenyl)methyl]-piperidine
15 1-(2-furanylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(2,6-dichlorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(4-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(3-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-(3-furanylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
20 1-[(4-ethylphenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(2-methylphenyl)methyl]-piperidine
1-[(3-chlorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
4-(1*H*-imidazol-4-ylmethoxy)-1-[(5-methyl-2-thienyl)methyl]-piperidine
1-[(4-bromo-2-thienyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
25 1-([2,2'-bithiophen]-5-ylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
3,5-dichloro-2-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-phenol
1-[(3,4-difluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(3,5-difluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
1-[(6-chloro-1,3-benzodioxol-5-yl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine

- 1-[[4-[4-(1,1-dimethylethyl)-2-thiazolyl]phenyl]methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
- 4-(1*H*-imidazol-4-ylmethoxy)-1-[(1-methyl-1*H*-pyrrol-2-yl)methyl]-piperidine
- 1*H*-indole, 3-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-1-(phenylmethyl)-
- 5 1-[(5-chloro-2-thienyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
- 1-(1,3-benzodioxol-4-ylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
- 2-thiophenecarbonitrile, 3-[[4-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]phenoxy]methyl]-piperidine
- 4-(1*H*-imidazol-4-ylmethoxy)-1-[[5-(phenylethynyl)-2-thienyl]methyl]-piperidine
- 10 4-(1*H*-imidazol-4-ylmethoxy)-1-[[5-(4-nitrophenyl)-2-furanyl]methyl]-piperidine
- 4-(1*H*-imidazol-4-ylmethoxy)-1-[[5-(3-nitrophenyl)-2-furanyl]methyl]-piperidine
- 1-[(4-chloro-1*H*-pyrazol-3-yl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
- 1-[(4-bromo-1-methyl-1*H*-pyrazol-3-yl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
- 1-[(4-bromo-1*H*-pyrazol-3-yl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
- 15 2-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-benzonitrile
- 4-(1*H*-imidazol-4-ylmethoxy)-1-[(4-iodophenyl)methyl]-piperidine
- 1-[(5-ethyl-2-thienyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
- 4-(1*H*-imidazol-4-ylmethoxy)-1-[[5-(methylthio)-2-thienyl]methyl]-piperidine
- 1-[[1-(3,5-dichlorophenyl)-1*H*-pyrrol-2-yl]methyl]-4-(1*H*-imidazol-4-ylmethoxy)-
- 20 piperidine
- 1-[[1-(4-chlorophenyl)-1*H*-pyrrol-2-yl]methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
- 4-(1*H*-imidazol-4-ylmethoxy)-1-[[4-(phenylethynyl)-2-thienyl]methyl]-piperidine
- 4-(1*H*-imidazol-4-ylmethoxy)-1-[(3-phenoxy-2-thienyl)methyl]-piperidine
- 1-[[2-chloro-5-(trifluoromethyl)phenyl]methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
- 25 4-(1*H*-imidazol-4-ylmethoxy)-1-[(4-propoxyphenyl)methyl]-piperidine
- 2-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-phenol
- 1-[(2,4-difluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
- 3-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-2-thiophenecarbonitrile
- 1-(benzo[*b*]thien-3-ylmethyl)-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
- 30 2-chloro-3-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-pyridine

3-[[4-(1*H*-imidazol-4-ylmethoxy)-1-piperidinyl]methyl]-2-(2-propenyl)-phenol
 1-[(4-chloro-3-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 4-(1*H*-imidazol-4-ylmethoxy)-1-[[4-(trifluoromethoxy)phenyl]methyl]-piperidine
 1-[(2,6-difluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-[(4-bromo-2-fluorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-[(2,2-difluoro-1,3-benzodioxol-5-yl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 1-[(4-butoxyphenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 4-(1*H*-imidazol-4-ylmethoxy)-1-[(2,3,5-trichlorophenyl)methyl]-piperidine
 1-[(2,5-dichlorophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 4-(1*H*-imidazol-4-ylmethoxy)-1-[[2-(trifluoromethyl)phenyl]methyl]-piperidine
 1-[(4-chloro-2-nitrophenyl)methyl]-4-(1*H*-imidazol-4-ylmethoxy)-piperidine
 and pharmaceutically acceptable salts and solvates thereof.

9. A compound of formula (IA):



in which:

Ar is an aryl group, a 5-7 membered heteraromatic ring containing 1-4 heteroatoms selected from nitrogen, oxygen or sulphur, or a bicyclic or tricyclic heteraromatic ring containing 1-4 heteroatoms selected from nitrogen, oxygen or sulphur, each of which can be optionally substituted by 1-3 groups selected from C₁₋₆ alkyl, C₁₋₆ alkylthio, C₁₋₆ alkoxy, halogen, cyano, CF₃, OCF₃, C₃₋₆ cyclolalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₂₋₆ alkenyloxy, hydroxyl, nitro, tosyl, thienyl, benzyl, phenyl, nitrophenyl,

R¹ is hydrogen or C₁₋₆ alkyl;

X is O, NR², CH₂ or SO_x

R^2 is C_{1-6} alkyl;

x is 0, 1 or 2;

5 Y is $C=O$, SO_2 , or $(C=O)NH$;

Z is $(CR^3R^4)_r$, or Y and Z together form a $CH=CH$ group;

m and n are independently 0, 1, 2 or 3;

10

p and q are independently 0, 1 or 2;

r is 0, 1, 2, 3, or 4, and

15 R^3 and R^4 are independently hydrogen or C_{1-6} alkyl.

10. A compound according to claim 9 in which Y is $C=O$.

11. A pharmaceutical composition comprising a compound of formula (IA), or a
20 pharmaceutically acceptable salt or solvate thereof, as claimed in claim 10 or 11 in
association with a pharmaceutically acceptable adjuvant, diluent or carrier.

12. A compound of formula (IA), or a pharmaceutically acceptable salt or solvate thereof,
as claimed in claim 10 or 11 for use in therapy.

25

13. Use of a compound of formula (IA), or a pharmaceutically acceptable salt or
solvate thereof, as claimed in claim 10 or 11 in the manufacture of a medicament for use in
therapy.

30

14. Use of a compound of formula (IA), or a pharmaceutically acceptable salt or
solvate thereof, as claimed in claim 10 or 11 in the manufacture of a medicament for use as
a histamine H_4 antagonist.

35

15. A compound of formula (IA) for use in the treatment of a disease where
modulation of histamine H_4 activity is beneficial.

ABSTRACT

5

The invention relates to piperidine compounds, processes and intermediates used in their preparation, pharmaceutical compositions containing them and their use in therapy.

PRV 03-07-21

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.